

Hatcher: Algebraic Topology

Kenny Workman

October 11, 2024

1 The Fundamental Group

1.1 Intuition

The Borromean rings are three linked circles where any two circles without a third are unlinked. It is a nice object that elucidates the difference between abelian and nonabelian fundamental groups.

Label the three rings A , B , and C . Consider C as an element of $\mathbb{R}^3 \setminus A \cup B$. C can be represented as $aba^{-1}b^{-1}$, where a and a^{-1} are forward and reverse oriented paths around A (and B). This element is not trivial, so the free group generated by a and b is not abelian.

Modify the ring so that A and B are linked. The same $C = aba^{-1}b^{-1}$ is now trivial. It is easiest to see this visually. Refer to the picture in Hatcher and slide the first loop a all the way around A until it is hanging over the circle. This loop is clearly contractible.

From this, we conclude there are two ways to tell if two circles are linked:

- Show that one of the circles is an element of the fundamental group of the complement of the other circle. (Simple example of loop wrapping around a circle.)
- Show the fundamental group of the complement of both of the circles is the abelian. (As in the Borromean ring example.)

1.2 Basic Constructions

Definition 1.1 (Path). A path in the space X is a continuous map $f : I \rightarrow X$ where I is the unit interval $[0, 1]$.

Definition 1.2 (Homotopy of paths). A continuous family of maps represented by the continuous map $f : I \times I \rightarrow X$

Example 1.3 (Linear homotopy). Consider two paths f, g that share endpoints ($f(0) = g(0) = x_0$ and $f(1) = g(1) = x_1$). Any such paths are always homotopic by the homotopy

$$h_t(s) = (1 - t)f(s) + (t)g(s).$$

Proposition 1.4 (homotopy of paths with fixed endpoints is an equivalence relation).

Definition 1.5 (The product of paths). For paths f, g where $f(1) = g(0)$, we can define the product $h = f \cdot g$ as

$$h(s) = \begin{cases} f(2s) & 0 \leq s \leq 0.5 \\ g(2s - 1) & 0.5 \leq s \leq 1 \end{cases}$$

Proposition 1.6 (The fundamental group is in fact a group). $\pi_1(X, x_0)$ with respect to the operator $[f] \cdot [g] = [f \cdot g]$ is a group.

Definition 1.7 (Reparameterization of a path). We can precompose any path f with $\phi : I \rightarrow I$ such that $f \circ \phi \simeq f$. $f \circ \phi$ is then the reparameterization of f .

Proof. To see our operator is well-defined, $[f \cdot g]$ depends only on $[f]$ and $[g]$. In other words, a homotopy exists between paths in $[f \cdot g]$ if and only if it is the composite of homotopies between paths in $[f]$ and $[g]$.

First, we verify associativity of the operator. Consider $(f \cdot g) \cdot h$. This product composes the composition of f and g with h . Visually, h is the first half and g and f occupy the last quarters of the product path. We can construct a piecewise function to change the proportions of these components - "shrink" h , keep g the same and "expand" f . $[(f \cdot g) \cdot h]\phi \simeq f \cdot (g \cdot h)$ gives us the desired equivalence.

Second, we check the identity property of the constant map c . f can be reparameterized by the piecewise function that speeds it up for the first half of I and holds it constant at $f(1)$ for the second half where $f\phi \simeq f \cdot c$.

Third, we verify the existence of an inverse for each $[f]$. Define $[\bar{f}]$ where $\bar{f}(s) = f(1 - s)$. Then $f \cdot \bar{f}$ is homotopic to the constant path. To see this construct $h_t = f_t g_t$ where $f_t = f$ on $[0, t]$ and $f_t = f(t)$ on $[1 - t, 1]$ and g_t is the inverse of this function. As t approaches 1, h_t has a larger constant region in its image starting from the middle and growing to the endpoints until $h_1(s) = f(0) = g(1)$ is our constant map at timepoint 1. □

Proposition 1.8 (The fundamental group is independent of the choice of basepoint up to isomorphism). *If f is a loop with basepoint x_1 and h is a path from x_0 to x_1 . The map $\beta_h : \pi_1(X, x_0) \rightarrow \pi_1(X, x_1)$ defined as $\beta_h([f]) = [h \cdot f \cdot \bar{h}]$ is an isomorphism.*

Proof. β_h is well-defined. To see this, if $[f] = [f']$ implies that f and f' are homotopic, so certainly $h \cdot f \cdot \bar{h}$ and $h \cdot f' \cdot \bar{h}$ are homotopic and $\beta_h([f]) = \beta([f'])$. β_h is a homomorphism. $\beta_h([f \cdot g]) = [h \cdot f \cdot g \cdot \bar{h}] = [h \cdot f \cdot h \cdot \bar{h} \cdot g \cdot \bar{h}] = [h \cdot f \cdot h] \cdot [h \cdot g \cdot \bar{h}] = \beta_h([f]) \cdot \beta_h([g])$. β_h is an isomorphism. □

Definition 1.9. A space is **simply connected** if it is path connected and has a trivial fundamental group.

Definition 1.10. lift Given $f : Y \rightarrow X$ and a covering space $p : \tilde{X} \rightarrow X$, a lift of f is a map $\tilde{f} : Y \rightarrow \tilde{X}$ that satisfies $f = p\tilde{f}$.

We are about to compute the fundamental group of the circle, but will use general properties of lifts (and covering spaces) to make the proof efficient.

(a) Given a path $f : I \rightarrow X$, and a covering space p , and some $\tilde{x}_0 \in p^{-1}(x_0)$, there exists a unique lifted path \tilde{f} starting at \tilde{x}_0 .

(b) Given a homotopy of paths in X , a covering space p , and some $\tilde{x}_0 \in p^{-1}(x_0)$, there exist a unique homotopy of lifted paths in \tilde{X} starting at \tilde{x}_0 .

We can see that both of these statements are specific versions of the more general statement (c), given a $F : Y \times I \rightarrow X$ and a $\tilde{F} : Y \times \{0\} \rightarrow \tilde{X}$ lifting $F|_{Y \times \{0\}}$, there exists a homotopy $\tilde{F} : Y \times I \rightarrow \tilde{X}$ that lifts F and restricts to the given $\tilde{F} : Y \times \{0\} \rightarrow \tilde{X}$. We'll prove (c) so we can use it later.

Proof. We construct a lift of an arbitrary neighborhood $N \times I$.

Consider a fixed $\{y\} \times I$ and notice that for each point (y, t) , $F((y, t))$ lies in some evenly covered open set U_t .

Because F is continuous, we can construct an open neighborhood $N_t \times (a_t, b_t)$ of y where $F(N_t \times (a_t, b_t)) \subseteq U_t$.

$\{N_t \times (a_t, b_t)\}_t$ cover $\{y\} \times I$. Since I is compact, we can recover a finite subcollection. Let N be the intersection of all the N_t in this subcollection and partition I into $\{[0, t_1] \cdots [t_i, t_{i+1}] \cdots [t_m, 1]\}$ in such a way where each closed interval lies within an open set in our subcollection. This partition need not be finite and probably is not if all of these closed intervals must lie entirely within the previously defined open sets.

Then $N \times [t_i, t_{i+1}]$ each contain (y, t) for some $t \in [t_i, t_{i+1}]$, and fixed y , while $F(N \times [t_i, t_{i+1}]) \subseteq U_i$, an evenly covered open set. Because each U_i is homeomorphic with some sheet \tilde{U}_i by p , Starting with $N \times [0, t_1]$, while we are only given \tilde{F} for $N \times \{0\}$, we can extend the map to the rest of this neighborhood by Fp^{-1} for p defined over the sheet that contains $\tilde{F}(N \times \{0\})$. We proceed by induction, defining \tilde{F} over $N \times [t_i, t_{i+1}]$ using Fp^{-1} over the same sheet \tilde{U}_i (t_i is shared by the last neighborhood and all neighborhoods must lie entirely in some sheet, so they both lie in the same sheet). The only additional adjustment might

be restricting N for $N \times [t_i, t_{i+1}]$ because it was only restricted sufficiently to lie within some U_i , but not the one homomomorphic to the sheet that we need (?). □

To see that (c) implies (a), consider a path as a homotopy of points. To see the same with (b), if we are given a homotopy of paths and the starting point of their lifted paths, the full lifted paths are uniquely determined by (a). If we consider Y as I , a homotopy of the lifted paths follows.

Theorem 1.11 ($\mathbb{Z} \cong \pi_1(S^1)$). *The fundamental group of S^1 is the infinite cyclical group generated by the loop $\omega(s) = (\cos 2\pi s, \sin 2\pi s)$*

Proof. First recognize that because $[\omega_n] = [\omega]^n$, where $\omega_n = z \mapsto (\cos 2\pi n z, \sin 2\pi n z)$, our proof is equivalent to showing any loop f is homotopic to some $[\omega_n]$ because it is then a product of our generator.

Let $p : \mathbb{R} \rightarrow S^1$ be a covering space defined as $p = s \mapsto (\cos 2\pi s, \sin 2\pi s)$. We will first state two useful properties of lifts with this covering space and use these to show our result without proof before proving them. □

We can immediately use $\pi_1(S^1)$ to prove important theorems. The big idea is that each full rotation around the circle is a unique element of the group. This suggests, among other things, that a loop once around cannot be continuously deformed to a loop twice around.

Our arguments proceed by contradiction, by assuming that our desired result does not hold and showing that this assumption implies some homotopy between loops on the circle that is not allowed.

Theorem 1.12 (Fundamental Theorem of Algebra). *Every non constant polynomial with coefficients in \mathbb{C} has at least one root in \mathbb{C} .*

Proof. Assume our polynomial $p(z) = z^n + a_1 z^{n-1} \dots a_n$ has no roots. Define the following $f_r : I \rightarrow \mathbb{C}$ for each real number:

$$f_r(s) = \frac{p(re^{2\pi i s}) \setminus p(r)}{|p(re^{2\pi i s})| \setminus |p(r)|}$$

We claim this is a loop in the unit circle S^1 . To see this, compute some values for fixed r and note for any value of r , $f_r(0) = 1$, $f_r(1) = 1$. Note that f_0 is a constant map with value 1 and is the trivial loop. Then by varying r we obtain a homotopy between any f_r and 0 with (the embedded) basepoint $(1, 0)$.

We now show that p must be constant. Choose a large r such that $r \geq \sum_n |a_n| \geq 1$. Observe then $r^n = r r^{n-1} \geq (\sum_n |a_n|) r^{n-1}$. This expression motivates a new homotopy for each f_r , let $f_{(r,t)}$ be our previous definition with $p_t(z, t) = z^n + t(z^{n-1} a_1 + \dots a_n)$ substituted for each p . See that f_t has no zeros on the circle of complex values with radius r satisfying our expression (this is why we constructed it at all). Note that $f_{(r,0)} = e^{2\pi i s t n}$ (check). Then $f_{(r,t)}$ is a homotopy between f_r and homotopy class $[w_n] \in \pi_1(S^1)$. But f_r is homotopic to 0. Then w_n is also homotopic to 0 and n must be 0. Our $p(z) = a_n$ must be constant. □

Theorem 1.13 (Brouwer fixed point theorem). *Any continuous map $h : D^2 \rightarrow D^2$ must have a fixed point $h(x) = x$.*

This theorem was initially proved by a gentleman named L.E.J. Brouwer circa 1910 and seems to be foundational to the rest of algebraic topology and other fields like differential topology.

We now introduce a theorem which proves that there must exist two antipodal locations on the surface of the earth with the same temperature and pressure (assuming that temperature and pressure are continuous functions of location).

Theorem 1.14 (Borsuk-Ulam theorem). *A continuous map $f : S^2 \rightarrow \mathbb{R}^2$ must have at least one pair of antipodal points x and $-x$ where $f(x) = f(-x)$.*

Lets build intuition for the two dimensional case by proving the theorem in one dimensions. For a map $f : S^1 \rightarrow \mathbb{R}$, construct continuous $h = f(x) - f(-x)$. Pick a point x , and evaluate h at points x and $-x$ halfway across the circle from each other. Notice $h(x) = -h(x)$ so there must exist some y where $h(y) = 0 \in [h(x), -h(x)]$ by intermediate value theorem. We proceed with proof in two dimensions:

Proof. Define a path $\eta(s) = (\cos 2\pi s, \sin 2\pi s, 0)$ around the equator of S^2 and a map $g : S^2 \rightarrow S^1$ defined as $g(x) = \frac{f(x)-f(-x)}{|f(x)-f(-x)|}$. Let $h = g\eta$. Check that η is nullhomotopic (one can shrink the equator to one of the poles) and therefore h is also nullhomotopic. (If $\eta \simeq 1$ by $\psi : I \times I \rightarrow S^2$, then $g\eta \simeq g1$ by $g\psi : I \times I \rightarrow S^1$)

See that because $g(-x) = -g(x)$ and $\eta(s + \frac{1}{2}) = -\eta(s)$, $g\eta(s + \frac{1}{2}) = g(-\eta(s)) = -g\eta(s)$, so $h(s + \frac{1}{2}) = -h(s)$. We now examine the lift \tilde{h} , induced by $p = s \mapsto (\cos 2\pi s, \sin 2\pi s)$, and see that $\tilde{h}(s + \frac{1}{2}) = \tilde{h}(s) + x$ where x are all the values that map $\tilde{h}(s) + x$ opposite to $\tilde{h}(s + \frac{1}{2})$ under the trigonometric functions composing p . Then $x = \frac{q}{2}$ where q is an odd integer. (Check this. Hatcher skips some steps that most readers would find obvious probably).

Think about what we are doing here - lifting h from its path on S^1 to \mathbb{R} - stretching it out "flat" so we can see how much it winds along S^1 .

If $\tilde{h}(s + 1/2) = \tilde{h}(s) + \frac{q}{2}$, we can solve for q , $q = 2(\tilde{h}(s + 1/2) - \tilde{h}(s))$, and see it cannot be a continuous function of s unless it is constant because it is constrained to integer values. Then $\tilde{h}(1) = \tilde{h}(\frac{1}{2}) + \frac{q}{2} = \tilde{h}(0) + q$, so \tilde{h} is a path that travels a distance q on \mathbb{R} , so must wind around S^1 q times (is q times a generator in S^1). So h is not nullhomotopic because q is not 0.

But we saw that h , as the composition $g \cdot \eta$, is nullhomotopic. This is a contradiction. □

Reflecting on the structure of this proof for a moment, it is the existence of some continuous map $g : S^2 \rightarrow S^1$. Intuition tells us that the nullhomotopy sliding the equator to a pole should do nothing when embedded in S^1

This theorem says a few things. It tells us the surface of a sphere can never be one-to-one with an embedding in R^2 , as two of the points always collapse, so it cannot be homeomorphic with a subspace of R^2 . This is obvious when you think of the stereographic projection of the sphere onto the plane, but is hard to prove in general.

Exercise 1.1 (8). Does the Borsuk-Ulam Theorem hold for the torus? For some continuous $f : S^1 \times S^1 \rightarrow \mathbb{R}^2$, does there always exist (x, y) and $(-x, -y)$ where $f((x, y)) = f((-x, -y))$.




Figure 1: A refresher on the meridian and longitude of the torus.

Proof. We can't do the exact same thing as the proof for S^2 because loops on the surface of the torus are not nullhomotopic.

Consider the stereographic projection p of the torus resting on the $0XY$ plane to this plane. This map is continuous. Each point, denoted $(x, y) \in S^1 \times S^1$ on the torus that share an image on the plane must lie on the same meridian circle, so must share the first coordinate. Then there cannot be antipodal points that satisfy $p((x, y)) = p((-x, -y))$. ($x = 0$ does not lie on S^1). □

Note (Explicit parameterization of torus). The above problem is mostly an exercise in visualizing geometry and explicit equations are useful to understand the shape under study.

For $0 \leq \theta \leq 2\pi, 0 \leq \phi \leq 2\pi$ and r, R constants (the radius of the meridian tube and distance from origin to center of the meridian tube respectively)

$$z = r \sin \theta$$

$$y = (R + r \cos \theta) \sin \phi$$

$$x = (R + r \cos \theta) \cos \phi$$

See the below figure.

Then continuous $(x, y, z) \mapsto (x, y)$, always maps (x, y, z) and $(-x, -y, z)$ to different points.




Figure 2: Illustration of parameterization of torus

Theorem 1.15 (Fundamental group of a product space). $\pi_1(X \times Y) \cong \pi_1(X) \times \pi_1(Y)$ if X and Y are path connected.

Proof. Recall a map into a product space $f : Z \rightarrow X \times Y$ is continuous iff constituent maps $g : Z \rightarrow X$ and $h : Z \rightarrow Y$ are continuous. Then each loop $\omega : I \rightarrow X \times Y$ has equivalent pairs of loops $\omega_X : I \rightarrow X$ and $\omega_Y : I \rightarrow Y$. Similarly, homotopies of loops the product space have equivalent homotopies in the constituent spaces, which we can use to show that $\phi : \pi_1(X, Y) \rightarrow \pi_1(X) \times \pi_1(Y)$ defined as $[f] \mapsto ([g], [h])$ is a group homomorphism. \square

Exercise 1.2 (1.1.10). From the isomorphism $\pi_1(X \times Y, (x_0, y_0)) \cong \pi_1(X, x_0) \times \pi_1(Y, y_0)$, it follows that loops in $X \times \{y_0\}$ and $\{x_0\} \times Y$ represent commuting elements in $\pi_1(X \times Y, (x_0, y_0))$. Construct an explicit homotopy demonstrating this.

Proof. Consider arbitrary loops $(g, 1) : I \times I \rightarrow X \times \{y_0\}$ and $(1, h) : I \times I \rightarrow \{x_0\} \times Y$. Our goal is to construct the homotopy that shows $[(g, 1)] \cdot [(1, h)] = [(1, h)] \cdot [(g, 1)]$.

Denote our isomorphism as ϕ , then $\phi([(g, 1)] \cdot [(1, h)]) = ([g], [1]) \cdot ([1], [h])$. This element is equivalent to $[(g \cdot 1), [1 \cdot h]] = ([1], [h]) \cdot ([g], [1])$ by the homotopy $(F_X, F_Y) : (I \times I \rightarrow X, I \times I \rightarrow Y)$. Because $\phi([(1, h)] \cdot [(g, 1)]) = ([1], [h]) \cdot ([g], [1])$, $F_{X \times Y} : I \times (I \times I) \rightarrow X \times Y = (t, (s_g, s_h)) \mapsto (F_X((t, s_g)), F_Y((t, s_h)))$ is the explicit homotopy in $\pi(X \times Y, (x_0, y_0))$ (where t is some point in time I and s_g, s_h are points in the domain of loops g and h) that shows $[(g, 1)] \cdot [(1, h)] \simeq [(1, h)] \cdot [(g, 1)]$. This homotopy induces commutativity and can be constructed from arbitrary $(g, 1)$ and $(1, h)$. \square

Example 1.16 (Torus). The fundamental group of the torus can be thought of as a pair of integers. Formally, $\pi_1(S^1 \times S^1) \cong \mathbb{Z} \times \mathbb{Z}$.

Definition 1.17 (Induced homomorphism). The map $\varphi : X \rightarrow Y$ where $\varphi(x_0) = y_0$ induces a homomorphism $\varphi_* : \pi_1(X, x_0) \rightarrow \pi_1(Y, y_0)$ defined as $\varphi_*([f]) = [\varphi f]$

We can briefly check some properties of this homomorphism. Let φ, ψ be maps:

- $(\varphi\psi)_* = \varphi_*\psi_*$

- $1_* = 1$

The first comes from the associativity of the composition of maps ($(\varphi\phi)f = \varphi(\phi f)$). The second is straightforward from the definition. These properties of the induced homomorphism make the fundamental group a functor.

The induced homomorphism allows us to describe relationships between topological spaces as relationships between fundamental groups. Geometrical structure is informed by algebraic and vice-versa.

For instance, showing there exist continuous maps between a shape of interest and a shape that admits a trivial fundamental group is a useful method to compute the fundamental group of the original shape. Some examples follow.

Theorem 1.18. $\pi_1(S^n)$ is trivial if $n \geq 2$

Proof. We use Lemma 1.15 but omit its proof. Given a loop f with basepoint x_0 in X , and some decomposition of X into open A_α where $X = \bigcup_\alpha A_\alpha$, $x_0 \in A_\alpha$ and $A_\alpha \cap A_\beta$ is path connected. There exists a homotopy from f to a product of loops in each A_α .

Consider $S^n = A_1 \cap A_2$ where $A_1 = S^n - \{y\}$ and $A_2 = S^n - \{z\}$ for y, z antipodal on S^n . $A_1, A_2 \approx R^{n-1}$. Then every f in S^n is the product $g \cdot h$, where g and h are loops in A_1 and A_2 respectively. Let $\psi : A_1 \rightarrow \mathbb{R}^{n-1}, \phi : A_2 \rightarrow \mathbb{R}^{n-1}$ be homeomorphisms, and because $\pi_1(\mathbb{R}^{n-1})$ is trivial, $[g], [h]$ are nullhomotopic in $\pi_1(A_1), \pi_1(A_2)$ under ψ_*, ϕ_* . So $[f] = [g] \cdot [h] = 1$ for any f . \square

Exercise 1.3 (1.1.12). Show that every homomorphism $\phi : \pi_1(S^1) \rightarrow \pi_1(S^1)$ can be realized as the induced homomorphism of the map $\varphi : S^1 \rightarrow S^1$.

Proof. We can consider the set of homomorphisms $\{\phi : \pi_1(S^1) \rightarrow \pi_1(S^1)\}$ as the equivalent set of homomorphisms $\{\phi : \mathbb{Z} \rightarrow \mathbb{Z}\}$.

First we see that the set of homomorphisms $\{\phi : \mathbb{Z} \rightarrow \mathbb{Z}\}$ are described by the set of maps $\{\phi' = x \mapsto mx \mid m \in \mathbb{Z}\}$. This set includes the identity and 0 map, but also all homomorphisms that scale group elements by an integer factor.

We then claim the set of maps $\{f_m : S^1 \rightarrow S^1 \mid m \in \mathbb{Z}\}$ where $f_m = (x, y) \mapsto (mx, my)$ can be realized as $\{\phi'\}$ as induced homomorphisms.

To see this, consider a loop in S^1 , ω_n , where n denotes the number of turns around the circle (formally $s \mapsto (\cos 2\pi ns, \sin 2\pi ns)$) and examine its transformation under f_m . f_m stretches (or shrinks) ω_n to a loop ω_{mn} . This is clear by looking at the lifts of ω_n and $f_m\omega_n$, which are maps over $\mathbb{R} \rightarrow S^1$ defined as $\omega_{n*} = x \mapsto (\cos 2\pi x, \sin 2\pi x)$ and $f_m\omega_{n*} = x \mapsto (\cos m2\pi x, \sin m2\pi x)$, to see that f_m indeed stretches (or shrinks) a " n turn" loop to an " $n * m$ " turn loop. Then $f_{m*} = [\omega_n] \mapsto [\omega_{n*m}]$ corresponds to a $\phi' : x \mapsto mx$ as desired and $\{f_m\}$ and $\{\phi'\}$ are in one-to-one correspondence. \square

Exercise 1.4 (1.1.13). Consider a space X and a path connected subspace A with basepoint $x_0 \in A \subseteq X$. Show that every map $\pi_1(A) \hookrightarrow \pi_1(X)$ is surjective iff each path in X that has its endpoints in A is homotopic to a path in A .

Proof. We begin with the reverse direction. A loop g in X with basepoint x_0 is certainly a path in X with its endpoints in A , so it is homotopic to a path in A with the same endpoints, a loop, f in A . Then for every $\theta : \pi_1(A) \hookrightarrow \pi_1(X)$, $[g] \in \pi_1(X, x_0)$ must be equal to some $\theta([f])$ as $f \simeq g$.

To see the forward direction, define h as the path in X with endpoints in A . We can turn h into a loop with the product $\eta h \bar{\mu}$ where η, μ are the paths from x_0 to the first and second endpoint respectively. Recall A is path connected so we can always construct such paths. Because all $\pi_1(A) \hookrightarrow \pi_1(X)$ are surjective, $\eta h \bar{\mu} \simeq h'$, where h' is a loop entirely contained in A . If we restrict the homotopy $F_t : I \rightarrow X$ to the closed subset of I that is h at time 0, the restricted homotopy shrinks h to a path in A , as desired. \square

Proposition 1.19. If X retracts to A , then i_* for $i : A \xrightarrow{X}$ is injective. If X deformation retracts to A , then i is an isomorphism.

Proof. If X retracts to A , $ri = 1$ (where r is the retraction). Because $(ri)_* = r_*i_* = 1_*$, i_* must be injective. If X deformation retracts to A , there exists some homotopy F_t where $F_t|_A = 1$ and $F_0 = 1$. Then any loop g in X with basepoint in A homotopes to some loop in A and $[g] = i_*([f])$ for some f . \square

So spaces that are deformation retracts have isomorphic fundamental groups.

Note that deformation retracts are rather strict examples of homotopies as not only do they fix a basepoint between X and A for all timepoints, but they restrict to the identity map over their retracting space for all I ($r_t|_{A \times I} = \mathbb{1}$).

Lets work towards an induced isomorphism under the more general condition of homotopy equivalence.

Consider a homotopy $\varphi_t : X \times I \rightarrow Y$ that is not a deformation retract but fixes the basepoint in X ($\varphi_t(x_0) = y_0$ for all t). Then $\varphi_{0*} = \varphi_{1*}$ as $[\varphi_0 f] = [\varphi_1 f]$. Any pair of induced homomorphisms under a basepoint preserving homotopy are equivalent.

Now consider homotopy equivalences that are also basepoint preserving. Let $\varphi : X \rightarrow Y$ and $\psi : Y \rightarrow X$ be such equivalences. Then $\psi\varphi \simeq \mathbb{1}$ but we just saw then $(\psi\varphi)_* = \mathbb{1}_*$ so φ_* is injective. φ_* is also surjective as $(\varphi\psi)_* = \mathbb{1}_*$.

We can actually see this is true even if our homotopy equivalence does not fix our basepoint. First lets show that induced homomorphisms are invariant under arbitrary homotopy when conjugated by path of basepoints.

Lemma 1.20. *Let $\varphi_t : X \rightarrow Y$ be a homotopy and $h|_{x_0 \times I}$ be a path between $\varphi_0(x_0)$ and $\varphi_1(x_0)$. Then $\varphi_{0*} = \beta_h \varphi_{1*}$*

Note. To see this, for a given loop f in X , we can build a new homotopy $h_t \cdot \varphi_t f \cdot \bar{h}_t$ of loops at $\varphi_0(x_0)$. Then by fixing timepoints 0 and 1, $\beta_h(\varphi_{0*}([f])) = \varphi_{1*}([f])$.

Now lets use this to show the pair of composition of maps, given by both directions of the homotopy equivalence, are isomorphic and then each constituent map is isomorphic.

Theorem 1.21 (The choice of basepoint between homotopy equivalent spaces is not important). *Consider the homotopy equivalence $\varphi : X \rightarrow Y$, then $\pi_1(X, x_0) \cong \pi_1(Y, \varphi(x_0))$.*

Proof. Consider the homotopy inverse $\psi : Y \rightarrow X$. Then $\psi\varphi \simeq \mathbb{1}$. We just saw that $(\psi\varphi)_* = \beta_h(\mathbb{1})_*$ for some path h from $\psi\varphi(x_0)$ to x_0 . Then $\psi_*\varphi_* = \beta_h$ is an isomorphism and φ_* is injective. Using the same argument, $\varphi\psi \simeq \mathbb{1}$ and φ is also surjective. \square

Exercise 1.5 (1.1.16). Show that a retraction $r : X \rightarrow A$ cannot exist for the following cases:

- $X = \mathbb{R}^3$ and A homeomorphic to S^1
- $X = S^1 \times D^2$ and A its boundary torus $S^1 \times S^1$
- $X = S^1 \times D^2$ and A is the circle shown (page 39)
- $X = D^1 \vee D^1$ and A is its boundary $S^1 \vee S^1$
- X is the disk with two opposite points identified and A is the boundary $S^1 \vee S^1$
- X is the mobius strip and A is its boundary circle.

Proof. • $\pi_1(A) \cong \mathbb{Z}$ and $\pi_1(X)$ is trivial. Then $i_* : \pi_1(A) \hookrightarrow \pi_1(X)$ cannot be injective.

- $\pi_1(A) \cong \mathbb{Z} \times \mathbb{Z}$ and $\pi_1(X) \cong \mathbb{Z}$. So i_* cannot be injective
- When A is embedded in the torus, it contracts to a point, so i_* is trivial. But $\pi_1(A) \cong \pi_1(S^1) \cong \mathbb{Z}$
- $\pi_1(A) \cong \mathbb{Z} \star \mathbb{Z}$ and $\pi_1(X) \cong \mathbb{1}$. i_* is cannot be injective.
- $\pi_1(X) \cong \mathbb{Z}$. To see this, there is a straightforward deformation retraction shrinking the disk to the x-axis. If our identified points are on the x-axis, this deformation retraction shrinks the disk to S^1 . $\pi_1(A) \cong \mathbb{Z} \star \mathbb{Z}$.
-

\square

Exercise 1.6 (1.1.18). Use Lemma 1.15 to show if a space X is obtained from attaching a e^n cell, where $n \geq 2$ to A , the induced inclusion $\pi_1(A) \hookrightarrow \pi_1(X)$ is surjective. Use this to show:

-
- For a path-connected CW complex X , the inclusion map of its 1-skeleton $X^1 \hookrightarrow X$ induces a surjection.

Proof. We know from Lemma 1.15 that we can express any loop f in X , with a basepoint $x_0 \in X \cap e^n$ (the boundary where the cell is glued to A), as a product $g \cdot h$, where g is in A and h is in e^n . Because X is path connected, if we have a loop in X , f' , with a basepoint x'_0 not on the boundary of the cell and A , we can construct a f with the desired basepoint x_0 that is on the boundary by conjugating f' with a path η from x'_0 to x_0 . Then $f = \bar{\eta} \cdot f' \cdot \eta$ and $f' \simeq f \simeq g \cdot h$. But $g \cdot h \simeq g$ because any h on e^n is contractible. Then any loop f is homotopic to a loop g in A , and because A is path connected, g is homotopic to some loop with a basepoint anywhere in A . This shows surjectivity on $\pi_1(A)$ with the map induced by $A \hookrightarrow X$.

- $\pi_1(S^1) \hookrightarrow \pi_1(S^1 \times S^2)$ is surjective by this result. It is also injective by mapping loops onto the S^1 component of the wedge sum. So it is an isomorphism and $\pi_1(S^1 \times S^2) \cong \mathbb{Z}$
- Choose some 1-cell on X and let X' be X where all of the n -cells attached to this boundary are removed.

□

Exercise 1.7 (1.1.3). $\pi_1(X)$ is abelian iff the basepoint-preserving homomorphism β_h depends only on the choice of endpoints of h

Proof. Consider homotopy classes $[f], [h]$ in X . By definition, $\beta_h([f]) = [hfh^{-1}]$. Consider a new constant path $c = s \mapsto h(0)$ (the basepoint of X). Then if $\beta_c = \beta_h$, $[f] = \beta_c([f]) = \beta_h([f]) = [hfh^{-1}] = [h][f][h^{-1}]$ for any choice of f, h . $\pi_1(X)$ is abelian.

To see the reverse, consider two arbitrary paths g, h (they need not be loops). Because $\pi_1(X)$ is abelian, $\beta_g([f]) = [g^{-1}fg] = [h^{-1}fh] = \beta_h([f])$ for any h .

□

Exercise 1.8 (1.1.5). Let X be some space. The following conditions are equivalent:

- Any map $S^1 \rightarrow X$ is homotopic to a constant map
- Any map $S^1 \rightarrow X$ extends to some map $D^2 \rightarrow X$
- $\pi_1(X, x_0) = 0$ for any $x_0 \in X$

Proof. To see (c) implies (a), consider any loop f . There exists a single homotopy class. Then f homotopic with the constant loop, $\mathbf{1}(I) = x_0$.

Now we show (a) implies (b). Let $f : S^1 \rightarrow X$ be our map and $h : S^1 \times I \rightarrow X$ be the homotopy relating f to some constant map. Let $g : D^2 \rightarrow S^1 \times I$ be a homeomorphism. Then hg is a continuous map where $hg|_{S^1} = f$.

Now we show (b) implies (c). Let $f : S^1 \rightarrow X$ be an arbitrary loop. Let $h : S^1 \times I \rightarrow D^2$ be a homotopy relating the identify on the circle to a constant map. Let g be the extension of f . Then gh homotopes f to a constant map whose image is a point in X . □

Exercise 1.9 (1.1.6). The canonical map $\phi : \pi(X, x_0) \rightarrow [S^1, X]$ is onto if X is path connected and places the conjugacy classes of X in one-to-one correspondence with $[S^1, X]$.

Proof. To see ϕ is surjective if X is path connected, consider $[f]$ in $[S^1, X]$. Let h be a path from x_0 to any point on the path f (note that h is constant if x_0 already lies on f). Then $\phi^{-1}([f]) = [hfh]$.

A proof of one-to-one correspondence between $[S^1, X]$ and conjugacy classes of π_1 is equivalent to showing $\phi([f]) = \phi([g])$ iff $[f]$ conjugates $[g]$ in π_1 .

To see the forward direction, let ψ_t be the basepoint free homotopy relating $\phi([f])$ and $\phi([g])$. Then $\psi_{0*} = \beta_h \psi_{1*}$, where h is the path induced by the image $\psi_t(0)$. Notice that h starts and ends at x_0 because

$[f]$ and $[g]$ had basepoints of x_0 under the preimage of ψ , so h is also a loop. Then $\psi_{0*} = \beta_h \psi_{1*}$ is equivalent to $[h \cdot g \cdot h] = [f]$ which implies $[h] \cdot [g] \cdot [h]^{-1} = [f]$.

To see the reverse direction, let $[h]$ be the element of π_1 that conjugates $[f]$ and $[g]$. □

1.3 Van Kampen's Theorem

To understand free products, let's review the direct products.

Definition 1.22 (Direct product of groups). Given an indexed set of subgroups $\{G_\alpha\}$, our direct product group $\prod_\alpha G_\alpha$ is the set of functions $\{f = \alpha \mapsto g_\alpha \in G_\alpha\}$.

Some facts that are useful to verify:

1. $\bigcap G_\alpha = 1$. (Every element is unique in the product, even if two copies of the same group are used in the product.)
2. For each factor group, there exists a surjective "projection homomorphism" $\pi_i : G \rightarrow G_\alpha$ where $G_\alpha = \{(1 \dots g_i \dots 1)\}$.
3. $G \setminus G_\alpha \cong \{(g_1, \dots, g_{\alpha-1}, g_{\alpha+1}, \dots, g_n)\}$ (To see this, construct a homomorphism that erases α component and invoke the First Isomorphism Theorem).
4. Elements of each factor group commute with each other.

Proposition 1.23 (Proof that direct products are commutative up to isomorphism.). *If $G = H \times K$, then $hk = kh$.*

Proof. Let $\psi : G \rightarrow H \times K$ be an isomorphism carrying elements of G to their tuple representation. $\psi(kh) = \psi(k)\psi(h) = (1, k)(h, 1) = (h, k) = \psi(hk)$. □

It is helpful to think of each element of the product as embedded in some latent tuple.

Now we might want to compose a group that is not commutative amongst factors. This will help us distinguish, for example, $\mathbb{Z} \star \mathbb{Z}$ (disjoint circles) from $\mathbb{Z} \times \mathbb{Z}$ (linked circles or torus). We introduce the free group:

To use the language of category theory, the free product is the coproduct in the category of groups.

Formally:

We show the binary operator of free groups is associative.

Associativity of free groups. □

The proof of Van Kampen's also relies on basic facts about homomorphisms and quotient groups. We derive some of these facts here to refresh the dome.

Proposition 1.24. *Consider $\phi : G \rightarrow H$ where $\ker \phi = K$. If $X = \phi^{-1}(a)$ is an element of G/K , then for any $u \in X$, $X = \{uk \mid k \in K\}$.*

. To see $uK \subseteq X$, observe $\phi(uk) = \phi(u)\phi(k) = \phi(u) \in X$. To see $X \subset uK$, consider $x \in X$ and notice that $\phi(u^{-1}x) = 1$. Then $u^{-1}x = k$ and $x = uk \subset uK$. □

Proposition 1.25. *Consider $\phi : G \rightarrow H$ where $\ker \phi = K$. The group operation in G/K is well-defined.*

. Consider arbitrary $X, Y \in G/K$ where $X = aK$ and $Y = bK$. Let $Z = XY$. To see multiplication is independent of choice of representative, pick arbitrary representatives $u \in X$ and $v \in Y$. Then $uv \in \phi^{-1}(ab)$. □

In fact, multiplication of quotient group elements is well defined under a more general condition - that our coset is a normal subgroup.

Proposition 1.26. *Let $uH, vH \in G/H$. Then $(uH)(vH) = uvH$ is well defined iff $H \trianglelefteq G$.*

. For the forward argument, choose arbitrary $x, x^{-1} \in G$. Then $(xH)(x^{-1}H) = H$, so $(xh)(x^{-1}1) \in H$. Then H is normal as desired. For the reverse argument, consider alternative representatives $u' \in uH$ and $v' \in vH$. We must show $u'v' \in uvH$. $u' = um$ and $v' = vn$ for some $m, n \in H$. Then $(um)(vn) = uvv^{-1}mvn$. But $v^{-1}mv \in H$ by assumption (denote this n'). Then $uvv^{-1}mvn = uvn'n \in uvH$. \square

Theorem 1.27 (Van Kampen's Theorem). *Let $i_{\alpha\beta} : \pi_1(A_\alpha \cap A_\beta) \rightarrow \pi_1(A_\alpha)$ be the homomorphism induced by the inclusion $A_\alpha \cap A_\beta \hookrightarrow A_\alpha$, $j_\alpha : \pi_1(A_\alpha) \rightarrow \pi_1(X)$ be the homomorphism induced by $A_\alpha \hookrightarrow X$, and $\phi : \star\pi_1(A_\alpha) \rightarrow \pi_1(X)$ be the homomorphism induced by each j_α .*

If each $A_\alpha \cap A_\beta$ is path connected, ϕ is surjective. If each $A_\alpha \cap A_\beta \cap A_\gamma$ is path connected, the kernel of ϕ is exactly the normal group generated by the elements $i_{\alpha\beta}(\omega)i_{\beta\alpha}(\omega)^{-1}$ for $\omega \in \pi_1(A_\alpha \cap A_\beta)$.

Proof. Observe $\phi(i_{\alpha\beta}(\omega)i_{\beta\alpha}(\omega)^{-1}) = j_\alpha i_{\alpha\beta}(\omega) j_\beta i_{\beta\alpha}(\omega)^{-1} = 1$ because $j_\alpha i_{\alpha\beta} = j_\beta i_{\beta\alpha}$. (Recall ϕ is the inclusion homomorphism applied to each letter in the word). Clearly all such elements of the free group are in the kernel of ϕ . We will show the normal group generated by such elements are exactly the kernel of ϕ .

To see N is the kernel of ϕ , we show that $\star_\alpha \pi_1(A_\alpha) \setminus N \cong \pi_1(X)$. Observe $[f_i]_\alpha N = [f_i]_\beta N$ by the definition of N ($[f_i]_\alpha [f_i]_\beta^{-1} \in N$). Furthermore $[f_1][f_i]_\alpha [f_2]N = [f_1][f_i]_\beta [f_2]N$ as N is normal so the group operation is well defined on cosets. \square

Note. We first look at wedge sums and use Van Kampen's to confirm our intuition - the fundamental group of the sum is the free product of the fundamental groups of summands.

Consider a wedge sum $\bigvee X_\alpha$ defined as the quotient space of the disjoint union $\coprod X_\alpha$ with basepoints identified. Let each X_α be path connected.

Then we can cover $\bigvee X_\alpha$ with $A_\alpha = U_\alpha \bigvee_{\beta \neq \alpha} X_\beta$.

Let $\phi : \star_\alpha \pi_1(A_\alpha) \cong \pi_1(\bigvee X_\alpha)$ be defined in the standard way.

Consider the intersection of two distinct covering sets, $A_\beta \cap A_\gamma$, and notice it is $\bigvee U_\alpha$, which is contractible. So ϕ is surjective. Furthermore, the intersection of three distinct covering sets is also contractible, so N is trivial.

Then ϕ is an isomorphism and the fundamental group of the wedge sum is the free product of the summands.

Exercise 1.10 (1.2.3). Show that the complement of a finite set of points in \mathbb{R}^n is simply connected if $n \geq 3$.

Proof. For each point x , let $A_x = \mathbb{R}^n - \{x\}$. This set is open and contractible, so its fundamental group is trivial. If X is the complement of the finite set of points in \mathbb{R}^n , then $\phi : \star\pi_1(A_x) \rightarrow \pi_1(X)$ is a surjective homomorphism by van Kampen's (Each $A_x \cap A_y$, which is \mathbb{R}^3 with two points removed, is certainly path-connected). Then $\phi^{-1}([f])$, for arbitrary $[f]$ in $\pi_1(X)$ is some word of loops in each A_x , each of which is nullhomotopic, so the entire word and therefore $[f]$ is nullhomotopic. ϕ is surjective, so $\pi_1(X)$ must be trivial and X is contractible. \square

Exercise 1.11 (1.2.4). Compute $\pi_1(\mathbb{R}^3 - X)$ where X is the set of n lines through the origin.

Proof. First see that $\mathbb{R}^3 - X \simeq S^2 \cap (\mathbb{R}^3 - X)$, by the homotopy $F_t = (x, t) \mapsto (1-t)x + t(\frac{x}{\|x\|})$. Note that $\mathbb{R}^3 - X$ has the origin removed so this map is well-defined.

$S^2 \cap (\mathbb{R}^3 - X)$ is now a sphere with $2n$ holes, a pair for each line through the origin. By the stereographic projection, $S^2 \cap (\mathbb{R}^3 - X) \simeq \mathbb{R}^2 - \{2n - 1 \text{ points}\}$. To see this, orient the sphere so that one of the origin passing lines is exactly the north-south pole. Under the projection, the pair of holes induced by this line collapse to the same hole, while the pairs of holes associated with all other lines project to distinct holes on \mathbb{R}^2 .

We can use Van Kampen's to compute $\mathbb{R}^2 - \{2n - 1 \text{ points}\}$ as the free group on $2n - 1$ generators. Decompose this space into $2n - 1$ open disks each with a single hole, denoted A_n . The intersection of any three, so also any pair, of sets is path-connected, so the $\star\pi_1(A_n)$ is isomorphic to $\pi_1(\mathbb{R}^2 - \{2n - 1 \text{ points}\})$, but the former is a free group on $2n - 1$ generators. \square

Note (Geometric interpretation of $\pi_1(\mathbb{R}^3 - X)$). Though we have computed $\pi_1(X)$ in the previous problem is the free group on $2n - 1$ generators, this is not very satisfactory visually. Consider X when two lines pass through the origin. It is not immediately clear why there must be three generators and not four visually.

Convince yourself that if we construct four loops, one around each "half" of the X and Y axis, that we can always construct one from some product of the other three (playing with orientation). See the image in [this post](#)




Figure 3:

Note (Stereographic projection of S^2). The explicit map $p : S^2 \rightarrow \mathbb{R}^2 - \{x\}, (x, y, z) \mapsto (\frac{2x}{1-z}, \frac{2y}{1-z})$, is derived from the algebraic relationship for each P and P' , $P' - N = t(P - N)$ with fixed and real t , and the requirement that P' lies on the XY plane. See the image above.

Work this out, good exercise.

Note (Interlude on quotient groups, relations, free groups). Consider the free group on two generators, $Fr(a, b)$, and impose the relation $a^3 = b^2$ (as in the torus knot $X_{3,2}$).

This same relation can be induced by "factoring out" the normal subgroup generated (conjugated by all elements in the group) by a^3b^{-2} . To see this, consider any word of the generators, such as $a^i b^j a^3 b^k$, and see for this word $a^i b^j a^3 b^k (b^{-k} a^{-3} b^k) = a^i b^j b^2 b^k$.

This idea is important in understanding abelianizations. The goal of the abelianization of a group is to produce the smallest possible group where elements commute, eg. $ab = ba$ for all $a, b \in G$. This is a relation, and as before, we can induce the relation by the quotient of some group generated by $aba^{-1}b^{-1}$. We just want to expand the generating subset to include all possible pairs of elements (so that they all commute) but this generating subset is the commutator subgroup, denoted $[G, G]$, given as $\{aba^{-1}b^{-1} \mid a, b \in G\}$. $G/[G, G]$ then gives a group with a new set of relations that are the commutativity properties for all possible pairs of elements.

(One thing that consistently tripped me up is the order of the elements in the normal group a^3b^{-2} . The leftmost letter should intuitively be an inverse, to "cancel out" and replace, eg. $a^3(a^{-3}b^2) = b^2$. But $a^{-3}(a^3b^{-2})a^3 = b^{-2}a^3$ is in our normal group as it is a conjugate. Similarly, generating our normal group with a^3b^{-2} (instead of $b^{-2}a^3$) produces an identical group. This is a simple idea but one worth documenting for precision.

Exercise 1.12 (1.2.10). Consider the arcs α and β embedded in $D^2 \times I$ as shown in Hatcher. A circle around the midpoint of the cylinder is nullhomotopic but show the same loop is not nullhomotopic in $D^2 \times I - (\alpha \cap \beta)$

Proof. Take one end of α and slide it towards the outside of the end of the cylinder, D^2 , and then along the outside of the side of the cylinder, until it reaches the opposite cylinder end. This new cylinder is the complement of a straightened line and a loop β , and is homotopic to the original cylinder. In fact, these

two spaces are homotopy equivalent, as composition of this initial sliding with the same sliding in reverse produces the original space, so they have isomorphic fundamental groups.

Now repeat this procedure with β and notice the final cylinder is the complement of two straightened lines. The fundamental group of this space is a free group on two generators. γ belongs to the homotopy class of a loop of two generators in the final space. This new space is in isomorphic correspondence with the original cylinder, so $[\gamma] \neq 0$, and we have our result. \square

Note. Group descriptions of basic objects

- $\pi_1(S^1) \cong \mathbb{Z}$
- $\pi_1(S^2) \cong 0$
- The fundamental group of a torus is $\pi_1(S^1 \times S^1) \cong \mathbb{Z} \star \mathbb{Z}$

We are now interested in how the fundamental group changes as we attach 2-cells to a cell complex.

Let X be some path connected space and Y be the result of attaching a collection of 2-cells $\{e_\alpha^2\}_\alpha$. The map $\varphi_\alpha : S^1 \rightarrow X$, corresponding to each such e_α^2 , defines where it is glued. The φ_α are loops that are not necessarily nullhomotopic in X but are certainly nullhomotopic when included in Y . Then φ_α lie in the kernel of the isomorphism $\pi_1(X, x_0) \hookrightarrow \pi_1(Y, y_0)$.

We will prove a stronger result, that the kernel of this isomorphism is exactly the normal group generated by these loops (conjugated by the path γ from a basepoint on φ to x_0). Let N be the normal subgroup generated by $\{\gamma_\alpha \varphi_\alpha \bar{\gamma}_\alpha\}_\alpha$.

Theorem 1.28 (van Kampen's applied to 2-dimensional cell complex). *(a) the kernel of the isomorphism $\pi_1(X, x_0) \hookrightarrow \pi_1(Y, y_0)$ is exactly N . So $\pi_1(X, x_0)/N \cong \pi_1(Y, y_0)$ (b) Consider Y obtained from X with the same procedure, but e^n for $n > 2$ used instead of e^2 , then N is trivial and $\pi_1(X, x_0) \cong \pi_1(Y, y_0)$. (c) For a path-connected cell complex X , the inclusion of its two-skeleton $X^2 \hookrightarrow X$ induces an isomorphism $\pi_1(X^2) \cong \pi_1(X)$.*

Proof. (a) We begin by constructing a space Z from Y , by attaching strips S_α along each γ . (See Hatcher for a precise description of this construction). What is more important is understanding why we do this, rather than proceeding with van Kampen's on Y itself.

Deconstruct Z into $A = Z - \cup_\alpha y_\alpha$ (where each y_α is a hole in the 2-cell e_α^2) and $B = Z - X$. Notice that B is contractible, as a space of many 2-cells glued together, so $\pi_1(B) = 0$. Verify quickly that A and B are open. B is a set of 2-cells, which are open, attached to some closed strips, so B is open. A is open as a two-skeleton with a finite number of holes.

When we setup the "van Kampen's homomorphism" in the standard way, $\pi_1(A) \star \pi_1(B) \rightarrow \pi_1(Z)$, we obtain the isomorphism $\pi_1(A) \star N \cong \pi_1(Z)$ where N is generated by the image of $\pi_1(A \cap B) \rightarrow \pi_1(A)$. (This is a straightforward application of van Kampen's theorem).

See that for each $\gamma_\alpha \varphi_\alpha \bar{\gamma}_\alpha$, we can obtain a corresponding $\delta_\alpha \in \pi_1(A \cap B)$ that travels along the strip and around some concentric circle in the corresponding e_α^2 . Indeed these loops are homotopic in $\pi_1(A \cap B)$ under a suitable basepoint change isomorphism.

It remains to show that the image of $\pi_1(A \cap B) \rightarrow \pi_1(A)$ is generated by these δ_α . We invoke van Kampen's again by decomposing $A \cap B$ into $A_\alpha = A \cap B - \cap_{\beta \neq \alpha} e_\beta^2$. This time, each such $A_\alpha \cap A_\beta$ is contractible, so N is trivial. Each A_α deformation retracts to φ_α , so $\pi_1(A_\alpha) \cong S^1$ or the group free on the generator δ_α (see this loop is homotopic to φ_α).

(b) To see this result, we proceed with the same decompositions as (a), but in the second invocation of van Kampen's we notice each A_α is now some n -cell with some hole, where $n > 2$, attached to many strips. $\pi_1(A_\alpha)$ is then trivial and so is N .

(c) When n is finite, the result follows from straightforward induction using (b). When n is infinite, the author is frankly uninterested in understanding the result at this time. \square

Theorem 1.29. *If $g \neq h$, then $\pi_1(M_g)$ and $\pi_1(M_h)$ are not isomorphic. Nor are the spaces homotopy equivalent.*

Proof. If $M_g \simeq M_h$, then their fundamental groups are isomorphic. These groups are the abelianizations of free groups in $2g$ and $2h$ generators respectively (this is immediate from the previous result). Because these groups are isomorphic, g and h must be the same. \square

Exercise 1.13 (1.2.14). Let X be the cube, $I \times I \times I$, where each face is identified with its opposite face with a half twist. Show this quotient space is the cell complex with 2 0-cells, 4 1-cell, 3 2-cells and 1 0-cell. Using this structure show that $\pi_1(X)$ is the quaternions group of order 8.

Proof. To first see the cell complex of X is composed of the described n-cells, we draw a picture identifying the appropriate vertices and edges. The 2-skeleton of the cube collapses to a pair of vertices with four edges between them. The 6 faces of the cube become 3 as we identify opposite pairs.

Recall the fundamental group of a cell complex can be described by its 2-skeleton as in 1.26. Then $\pi_1(X) \cong \pi(X^1)/N$, where X^1 is the 1-skeleton and N is the normal group generated by loops of the attached 2-cells as in 1.26. The single 3-cell has no effect on the fundamental group.

Calculating $\pi_1(X^1)$ is a straightforward application of van Kampen's. We decompose X^1 into three open sets A_1, A_2, A_3 , each obtained by tracing a different cycle through the two vertex graph X . If we assume X inherits a metric topology from \mathbb{R}^3 , these sets are open, and the intersection of any pair is a point. $\pi_1(A_1) \cong \pi_1(A_2) \cong \pi_1(A_3) \cong \mathbb{Z}$, so $\pi_1(X^1)$ is the group free on three generators.

To construct N , it is sufficient to trace the attachment of 2-cells to the front, side and top face of the cube. If we do this, using the hand drawn image as a guide, we obtain $\{abcd, adb^{-1}c^{-1}, b^{-1}dca^{-1}\}$ as the generators of N . $\pi_1(X)$ is then the free group on three generators with the relations induced by dividing by N :

$$\langle ab, ac^{-1}, ad \mid abcd = adb^{-1}c^{-1} = b^{-1}dca^{-1} = 1 \rangle$$

To see this group is isomorphic to Q_8 , let $i = ab, j = ac^{-1}, k = ad$ and rewrite the above presentation:

$$\langle i, j, k \mid ij^{-1}k = ki^{-1}j = i^{-1}kj^{-1} = 1 \rangle$$

We can rewrite the relations as $ij = k, i = jk, j = ki$ and then $k^2 = i^2 = ijk$ (from the first and second relation). Obtaining $j^2 = ijk$ comes from $j = k(i) = k(jk)$, so $j^2 = (jk)jk = ijk$.

$$\langle i, j, k \mid i^2 = j^2 = k^2 = ijk \rangle \cong Q_8 \cong \pi_1(X)$$

\square

Exercise 1.14. Show the fundamental group of the surface of infinite genus is free on infinite generators.

Proof. First we show that for $M_1 - \{x, y\}$, the torus with two holes, $\pi_1(M_1 - \{x, y\}) \cong \mathbb{Z} \star \mathbb{Z}$. To see this, consider the representation of a torus as a rectangle with opposite sides identified as in the figure. If we poke a hole in this shape, it deformation retracts to its 1-skeleton, which is the wedge sum $S^1 \vee S^1$. Poking an additional hole will not \square

We will use the fact that a punctured n-sphere is homeomorphic to the

Proposition 1.30 (Proof of generalized stereographic projection). *There exists a homeomorphism between $S^n - 1$ and*

Exercise 1.15 (1.2.4). Compute $\pi_1(\mathbb{R}^3 - X)$ where X is the union of n lines through the origin.

Proof. Observe the complement deformation retracts to S^2 with $2n$ points removed (the poles of our n lines). This object is homeomorphic to \mathbb{R}^2 with $2n - 1$ points removed. (maybe construct the explicit stereographic projection). We can use van Kampen's to show the fundamental group of this space is the free group of $2n - 1$ generators. \square

Exercise 1.16 (1.2.6). Suppose a space Y is obtained from a path-connected space X by attaching n -cells for a fixed $n \geq 3$. Then show the inclusion $X \hookrightarrow Y$ induces an isomorphism on π_1 . Use this result to then show that the complement of a discrete subspace of \mathbb{R}^n is simply-connected if $n \geq 3$.




Figure 4:

Proof. We proceed using the structure of the proof of 1.26. Let A be Y with a hole in each of the attached n -cells. Let B be the union of the attached n -cells. See that A and B are both open and $Y = A \cup B$, so we invoke van Kampen's theorem: $\pi_1(Y) = \pi_1(A) \star \pi_1(B)/N$ where N is defined in the usual way.

As before, A deformation retracts onto X , so $\pi_1(A) \cong \pi_1(X)$, and B is contractible, so $\pi_1(B) = 0$. Then it remains to show that N generated by $\{i_{AB}(\omega)i_{BA}(\omega)^{-1} \mid \omega \in \pi_1(A \cap B)\}$ is trivial (where the map i_{AB} is defined as $i_{AB} : \pi_1(A \cap B) \rightarrow \pi_1(A)$). See that any loop $i_{AB}(\omega)$ is contractible, because despite introducing a hole, the loop can homotope around it within the interior of the n -cell (if $n \geq 3$). Any $i_{BA}(\omega)$ is trivially contractible, being included into an open subset of an n -cell with no holes. Then $\pi_1(Y) \cong \pi_1(X)$.

To prove our last result, let X be our space of interest (the complement of some discrete subspace of R^n), we can obtain a covering $\{A_\alpha\}_\alpha$ where each $A_\alpha = X - x_\alpha$ is the complement of each point in the discrete subspace. See each A_α is open. Then $\pi_1(Z) = \star \pi_1(A_\alpha)/N$. □

Exercise 1.17 (1.2.7). Construct a cell complex for the two-sphere with north and south poles identified and use this to compute the fundamental group of this space.

Proof. We first construct the cell complex. Begin with a 0-cell x and attach a 1-cell e to x to form a circle. Then take a 2-disk σ and attach its boundary circle $\delta\sigma$ to e in two pieces, $\delta_1\sigma$ traces one endpoint to the other and attaches to e , while $\delta_2\sigma$ completes the circle in the same orientation but attaches to \bar{e} .

Then, by van Kampen's, our fundamental group is isomorphic to $\langle a \mid aa^{-1} \rangle$, but this is just the free group on one generator, so $\pi_1(X) \cong \mathbb{Z}$. □

Seeing that the cell complex is the same as the sphere with north and south poles identified was not immediately obvious.

This post was helpful (notation was taken from Lee Mosher's solution), viewing our space as the quotient map $f : D_2 \rightarrow X$ performing exactly two identifications. 2-disks can take different forms equivalent up to homeomorphism and their boundaries can simply be circles. Seeing that S^2 is simply D_2 attached along e and \bar{e} , where e is an arc connecting the poles, was crucial.

Exercise 1.18 (1.2.8). Consider a surface M_g of genus g that is split into two closed surfaces M_h and M_k by a circle C . M'_h and M'_k are constructed by deleting an open disk from each. Show that neither M'_h or M'_k retract to their boundary circle, but both (and all of M_g) retract to a lateral C' .

More familiarity with the two torus (really genus g surfaces in general) and practice with abelianization is needed to tackle this problem.

Note (Abelianization of $\pi_1(M_1)$). Consider $\pi_1(M_1) = \langle a, b \mid [a, b] \rangle$ as the familiar free group on two generators modulo the commutator (where $[a, b] = aba^{-1}b^{-1}$).

The abelianization of $\pi_1(M_1)$ is then $\pi_1(M_1) / [\pi_1(M_1), \pi_1(M_1)]$




Figure 5: The geometry of the two torus. Hilbert’s *Geometry and the Imagination*

1.4 Covering Spaces

Covering spaces let us (1) calculate fundamental groups of structures and (2) think about algebraic properties using geometric intuition. (I do not really understand how yet, so revisit this explanation after some exercise.)

Definition 1.31. Given some space X , the space \tilde{X} , along with map $p : \tilde{X} \rightarrow X$, is a **covering space** if each $x \in X$ has some open neighborhood U , where $p^{-1}(U)$ is a disjoint union of open sets and each one is mapped homeomorphically onto U by p . These open sets are called **sheets**.

$p^{-1}(U)$ is allowed to be empty, so p need not be surjective.

The helicoid (denoted S) is a good example that is easy to visualize. The helicoid can be parameterized as:

$$\begin{aligned} x &= s \cos 2\pi t \\ y &= s \sin 2\pi t \\ z &= t \end{aligned}$$

for $s \in (0, \infty), t \in \mathbb{R}$




Figure 6:

Then $p : S^1 \rightarrow \mathbb{R} \setminus \{0\}$ given by $(x, y, z) \mapsto (x, y)$ defines a covering space of $\mathbb{R} - \{0\}$.

I found this is best seen by looking at single line segments extending radially from the origin. Consider $X = (0.5, 1.5) \times \{0\}$. $p^{-1}(X)$ is a collection of disjoint segments. Our parameterization allows this segment of the x axis to exist whenever we make one full rotation around the circle. There are an infinite number of these segments and each maps to X homeomorphically.

We can also look at the covering spaces of S^1 . Let $p : S^1 \rightarrow S^1$ where $p(z) = z \mapsto z^n$ with $z \in \mathbb{Z}$. Here $z \in S^1$ is the complex number with $|z| = 1$, so $S^1 = \{z \in \mathbb{C} \mid |z| = 1\}$.

Let $U = (0, \frac{\pi}{2})$ be some open segment of the circle, if $n = 3$, what is $p^{-1}(U)$?

Once place to see the utility of covering spaces is with oriented graphs.




Figure 7:

Let X (Fig. 3) be basepoint with two loops labeled a and b . Now consider any graph, \tilde{X} , where edge vertex has four edges. If we label, with a or b , and orient each edge we obtain a structure that looks like X (respecting orientation!) as we zoom into each vertex. We can call this an **2-oriented graph**.

If we construct a map $p : \tilde{X} \rightarrow X$, where the interior of each edge in \tilde{X} maps homeomorphically to corresponding labeled edge in X following the orientation of both edges, we satisfy the properties for a covering space. It follows both that every oriented graph is a covering space of X and every covering space of X can be 2-oriented (every vertex has 4 edges that can be labeled so the local picture looks like X).

We are given a table of some of these \tilde{X} .

Theorem 1.32. $\pi_1(X) \cong \langle a, b^2, bab^{-1} \rangle$

Proof. Hatcher instructs us that we can use Van Kampen's. □

When we consider the fundamental group of the covering space as the "image subgroup" of $\pi_1(X)$, $p_*(\pi_1(\tilde{X})) \leq \pi_1(X)$, we recover yet another beautiful correspondence between algebra and geometry - there is a one to one correspondence between the subgroups $\pi_1(X)$ and covering spaces of X .

A few more algebraic interpretations of covering spaces:

- If we change our basepoint in a covering space that is otherwise the same, $p_*(\pi_1(\tilde{X}, x_1))$ is a conjugacy subgroup of $p_*(\pi_1(\tilde{X}, x_0))$ in $\pi_1(X)$ where the conjugating element is the loop connecting the basepoints.
- If a **symmetry** is an automorphism on a graph that preserves labels and orientations, then a graph can be "more" symmetric if it has more automorphisms and "the most" symmetric if every possible permutation of the graph vertexes have automorphisms with the desired properties. We will show that "the most symmetric" graphs are exactly those induced subgroups that are normal in $\pi_1(X)$.

We now define three **lifting properties** of covering spaces and show some applications. Given some covering space and a map into X , these properties tell us when a lift of a homotopy exists, when a lift of a regular map exists and when lifts are unique.

Proposition 1.33 (Homotopy lifting property). *Given a covering space $p : \tilde{X} \rightarrow X$, a homotopy $f_t : Y \rightarrow X$ and a lift of timepoint 0 of the homotopy $\tilde{f}_0 : Y \rightarrow \tilde{X}$, there exists a unique lift of the entire homotopy $\tilde{f}_t : Y \rightarrow \tilde{X}$.*

When Y is a point, this tells us that paths in X induce unique paths in the covering space.

When Y is I , this tells us that homotopies in X induce unique homotopies in the covering space.

The key idea here is that these homotopies are unique. If we consider the lift of a constant path, this implies that the lifted path is also constant

Note. Check this. If $p\tilde{f} = f$ and f is constant, if \tilde{f} had anything other than $\tilde{f}(0)$ in its image, what does that imply about the set of allowed \tilde{f} s? If \tilde{f} is itself constant, then $p\tilde{f} = f$ obviously holds.

We will use this to show p_* is injective.

Proposition 1.34. *The induced homomorphism $p_* : \pi_1(\tilde{X}, \tilde{x}_0) \rightarrow \pi_1(X, x_0)$ is injective. The image subgroup $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ consists of loops with basepoint x_0 that lift to loops in \tilde{X} with basepoint \tilde{x}_0 .*

Proof. To prove injectivity, consider two distinct homotopy classes, $[\tilde{f}'_0]$ and $[\tilde{f}''_0]$, that have the same image under p_* , so $p_*([\tilde{f}'_0]) = p_*([\tilde{f}''_0])$. Indeed $p\tilde{f}'_0 \simeq p\tilde{f}''_0$, given by some homotopy f_t , so our homotopy lifting property gives us a lifted homotopy \tilde{f}_t where $\tilde{f}'_0 \simeq \tilde{f}''_0$. Then $[\tilde{f}'_0] = [\tilde{f}''_0]$, so p_* is injective.

(This is a bit more explicit than Hatcher's proof using the kernel of p_*)

A loop in X with basepoint x_0 that lifts to a loop in \tilde{X} with basepoint \tilde{x}_0 is by definition an element of $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$.

The second statement is a bit bizarre, because should every element in the image subgroup, represented as $p_*\tilde{f}$, admit a straightforward lift \tilde{f} . I do not see where I have to invoke homotopy lifting, unless you consider some other representative of each homotopy class $g \simeq p\tilde{f}$, then $\tilde{g} \simeq \tilde{f}$, so the representative admits the desired lift. \square

Proposition 1.35. *The number of sheets in a covering space p is equal to the index of $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ in $\pi_1(X, x_0)$*

Proof. Let g be a loop in $\pi_1(X, x_0)$. \tilde{g} is its lift by the homotopy lifting property. Note that \tilde{g} need not be a loop as $[g] \notin p_*(\pi_1(\tilde{X}, \tilde{x}_0))$! (Consider the lift of a single wrap around the circle in $p : \mathbb{R} \rightarrow S^1$.)

Each representative of our coset $H = p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ can be denoted $H[g] = h \cdot g$, which has a lift $\tilde{h} \cdot \tilde{g}$. \tilde{h} is a loop at \tilde{x} by the prior theorem, so notice that $\tilde{h} \cdot \tilde{g}$ is a path that starts at \tilde{x}_0 and ends at $\tilde{g}(1)$.

We claim that these cosets are one to one with the preimage of basepoints $p^{-1}(x_0)$. Consider the function $\phi = [H]g \mapsto \tilde{g}(1)$. It is well-defined as for $hg, h'g \in [H]g$, $\phi(hg) = \phi(h'g) = g(1)$.

\tilde{X} is path-connected so the path \tilde{g} can be extended to any point in $p^{-1}(x_0)$, so ϕ is surjective. To see ϕ is injective, if $\phi([H]g) = \phi([H]g')$, then $g(1) = g'(1)$, so $g \cdot \bar{g}'$ is a loop with basepoint x_0 that lifts to a loop with basepoint \tilde{x}_0 . Then $g \cdot \bar{g}' \in H$ and $[H]g = [H]g'$. \square

We have two more properties. One will show when a lift exists and another will show when lifts are unique.

Proposition 1.36 (Lifting criterion). *Given a covering space $p : \tilde{X} \rightarrow X$ and some map $f : Y \rightarrow X$, there exists a lift of this map, denoted $\tilde{f} : Y \rightarrow \tilde{X}$, iff $f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$.*

Proof. The necessary condition is straightforward. The existence of a lift implies that $f = p\tilde{f}$, then $f_* = p_*\tilde{f}_*$. Then for any loop $\gamma : I \rightarrow Y$ in $\pi_1(Y, y_0)$, $p_*^{-1}f_*(\gamma) = \tilde{f}_*(\gamma)$. Certainly $f_*(\pi_1(Y, y_0)) \subseteq p_*(\pi_1(\tilde{X}, \tilde{x}_0))$.

The sufficient condition is less straightforward. We proceed by constructing our lift \tilde{f} explicitly. Let $\tilde{f} = y \mapsto \tilde{f}\gamma(1)$, where γ is any path from y_0 to y . We will show that this function is well-defined - that it does not matter what path we draw to y .

Consider another path γ' . We build a loop $\gamma \cdot \bar{\gamma}'$. Consider $f(\gamma \cdot \bar{\gamma}')$, which is also a loop. By assumption, this is some element of $p_*(\pi_1(\tilde{X}, \tilde{x}_0))$, call it $[x]$, so $f(\gamma \cdot \bar{\gamma}')$ homotopes to x . Because x lifts to a loop with basepoint \tilde{x}_0 , by the homotopy lifting property, $f(\gamma \cdot \bar{\gamma}')$ itself lifts to a loop with basepoint \tilde{x}_0 . So for $\tilde{f}(\gamma \cdot \bar{\gamma}')$, $\tilde{f}(\gamma)(1) = \tilde{f}(\bar{\gamma}')(0)$ and $\tilde{f}(\gamma)(1) = \tilde{f}(\gamma')(1)$ as desired.

We must also check for continuity. Consider some open U about $f(y)$ that is mapped homeomorphically to its lift \tilde{U} about $\tilde{f}(y)$ by $p : \tilde{U} \rightarrow U$. By continuity of f , there exists some N about y where $f(N) \subset U$. We will see that $\tilde{f}|_N = p^{-1}f$. N is path connected, so let η to be the path from y to any $y' \in N$ (γ is still the path from y_0 to y), then $\tilde{f}(N) = p^{-1}f(N)$ (Varying η , $\tilde{f} \cdot \eta(1)$ are exactly the points mapped onto \tilde{N} by p^{-1} . This is true for arbitrary N . \square)

Proposition 1.37 (Unique lifting property). *Given a covering space $p : \tilde{X} \rightarrow X$ and some map $f : Y \rightarrow X$, if \tilde{f}_1 and \tilde{f}_2 agree on one point of Y and Y is connected, they agree on all of Y .*

Proof. For a point $y \in Y$ let U be an evenly-covered neighborhood of $f(y)$ (recall this means that $p^{-1}(U)$ is a set of open sets in \tilde{X} each mapping homeomorphically to U). Let \tilde{U}_1 and \tilde{U}_2 be the open sets that $\tilde{f}_1(y)$ and $\tilde{f}_2(y)$ reside in respectively.

If $\tilde{f}_1(y) \neq \tilde{f}_2(y)$, then $\tilde{U}_1 \neq \tilde{U}_2$ so these sheets are disjoint.

Consider an open N about y that maps into \tilde{U}_1 and \tilde{U}_2 by each lift. Such an N exists because of continuity. If $\tilde{f}_1(y) = \tilde{f}_2(y)$, then N must map into the same \tilde{U} . Because $p\tilde{f}_1 = p\tilde{f}_2$ and p is injective over \tilde{U} (it is homeomorphic with U), these lifts agree over N . \square

Definition 1.38 (semilocal simple connectedness). For each point $x \in X$, there exists U , where $\pi_1(U) \hookrightarrow \pi_1(X)$ induced by the inclusion map is trivial.

Notice that this does not mean that U itself is simply connected, rather that loops within U can homotope throughout the whole space to X .

For a fixed, path-connected X , we are now interested in classifying the various covering spaces. The main result is a correspondence between subgroups of $\pi_1(X, x_0)$ and covering spaces $p : \tilde{X} \rightarrow X$, where $p_*(\pi_1(\tilde{X}))$ is equivalent to each subgroup. We are first interested in constructing the trivial subgroup for arbitrary X . Since we have shown that p_* is always injective, it is sufficient to construct a simply-connected covering space \tilde{X} .

Example 1.39 (Constructing the universal cover). What follows is a rather abstract procedure for building this simply-connected space, called the **universal cover**, for a path-connected, locally path-connected and semilocally simply-connected space X .

First, recall that there is a single homotopy class of paths for fixed endpoints in a simply connected space. If \tilde{X} is simply-connected and we fix some basepoint \tilde{x}_0 , then for each point $x \in \tilde{X}$, a path γ from x_0 to X uniquely defines $[\gamma]$. The set of $[\gamma]$ is bijective with \tilde{X} . By the homotopy lifting property, homotopy classes of paths starting at x_0 in X lift uniquely to homotopy classes of paths in \tilde{X} starting at \tilde{x}_0 . This gives a way to build \tilde{X} from just X . (But this does not mean that $\gamma(1)$ uniquely defines each point.)

$$\tilde{X} = \{[\gamma] \mid \gamma \text{ is a path in } X \text{ starting at } x_0\}$$

Let $p = [\gamma] \mapsto \gamma(1)$ be our covering map. p is surjective because X is path connected. We can have many distinct $[\gamma]$, all with the same endpoint, so p is certainly not injective (but most covering spaces are not so this is fine).

Now we make use of the semilocally simply-connected property. Let \mathcal{U} be the collection of U where $\pi_1(U) \rightarrow \pi_1(X)$ is trivial. Notice that \mathcal{U} is a basis for the topology of X . Every $x \in X$ has some U about it that satisfies this property and for each each pair of $U_1, U_2 \in \mathcal{U}$ and $x \in U_1 \cap U_2$, there exists $U_3 \in \mathcal{U}$ where $x \in U_3 \subseteq U_1 \cap U_2$ ($\pi_1(U_3) \rightarrow \pi_1(U_1 \cap U_2) \rightarrow \pi_1(U_1) \rightarrow \pi(X)$ is a chain of trivial homomorphisms induced by inclusions)

We need to now define a topology on \tilde{X} . Then show that p has the properties of a covering map (continuous + disjoint sheets). We define:

$$U_{[\gamma]} = \{[\gamma \cdot \eta] \mid \eta \text{ is a path within } U \text{ where } \eta(0) = \gamma(1)\}$$

See that restricting p to each U_γ now gives us a bijection. Surjectivity, as before, comes from U being path-connected. But now, if $[\gamma'] \neq [\gamma'']$, because $\pi_1(U)$ is trivial under inclusion, these paths must have different endpoints. So p is injective.

To prove one of the properties of a basis, we show in general that $U_{[\gamma]} = U_{[\gamma']}$ if $[\gamma'] \in U_{[\gamma]}$ (Certainly $U_{[\gamma']}$ $\subseteq U_{[\gamma]}$ as $[\gamma \cdot \eta'] = [\gamma \cdot \eta]$). Similarly $U_{[\gamma]} \subseteq U_{[\gamma']}$ as $[\gamma\eta] = [\gamma \cdot \eta \cdot \bar{\eta} \cdot \eta']$).

Then, varying over $U \in \mathcal{U}$ and $[\gamma]$ that induce distinct $U_{[\gamma]}$ for fixed U , $\{U_{[\gamma]}\}$ defines our topology on \tilde{X} . Every $[\gamma]$ has such a set. For $[\gamma''] \in U_{[\gamma]} \cap V_{[\gamma']}$, let $W \subseteq U \cap V$ contain $\gamma''(1)$. Then $[\gamma''] \in W_{[\gamma'']} \subseteq U_{[\gamma]} \cap V_{[\gamma']}$.

We now show that $p : U_{[\gamma]} \rightarrow U$ is also a homeomorphism. In other words, open $V \subseteq U$ are one-to-one with $V_{[\gamma]}$. Indeed, $p(V_{[\gamma]}) = V$. Similarly, $p^{-1}(V) \cap U_{[\gamma]} = V_{[\gamma]}$.

This is what we need to prove that p is a covering space. p is continuous, as we have showed p to be a homeomorphism when restricted to each element of the basis for \tilde{X} . To see $p^{-1}(U)$ is a set of disjoint sheets, see that $U_{[\gamma]}$ partitions this set as we vary γ . Indeed if $[\gamma''] \in U_{[\gamma]} \cap U_{[\gamma']}$, then $U_{[\gamma]} = U_{[\gamma'']} = U_{[\gamma']}$. (Stop and think about what disjoint sheets imply about γ for fixed U).

Finally we show that \tilde{X} is simply-connected. See first that for some path γ in X , a lift of this path in \tilde{X} can be described by the function $t \mapsto [\gamma_t]$ where γ_t is γ during $[0, t]$ but constant during $[t, 1]$. Now we show that $p_*(\pi_1(\tilde{X}), \tilde{x}_0)$ is trivial, and because p_* is injective, this gives us our result. Consider an element $g \in p_*(\pi_1(\tilde{X}))$, a loop at x_0 that lifts to a loop in \tilde{X} . This lift goes from $[x_0]$ to $[g_1]$, using the definition at the beginning of the paragraph, but $[g_1] = [x_0]$ because it is a loop. This is true for arbitrary $[g_1] = [g]$.

Example 1.40. Let $X_{m,n}$ be the quotient space of a cylinder $S^1 \times I$ with the identifications $(z, 0) \simeq (e^{\frac{2\pi i}{m}} z, 0)$ and $(z, 1) \simeq (e^{\frac{2\pi i}{n}} z, 1)$. Note z is a complex coordinate and the transformation is a $\frac{1}{n}$ twist.

A and B are subspaces of X and the quotients of $S^1 \times [0, \frac{1}{2}]$ and $S^1 \times [\frac{1}{2}, 1]$ respectively. One end of each half cylinder is just the circle ($A \cap B = S^1$), but they "twist" to the identification.

See that if $m = n = 2$, then A and B are the mobius band.

The procedure used to build the universal cover can be used to complete the rest of the subgroups.

Theorem 1.41. For each $H \leq \pi_1(X, x_0)$, there exists a covering space $p_* : X_H \rightarrow X$ where $p_*(\pi_1(X_H, \tilde{x}_0)) = H$ for a suitably chosen basepoint \tilde{x}_0 .

Proof. Our strategy is to construct a quotient space out of our universal cover.

For $[\gamma], [\gamma'] \in \tilde{X}$ (remember that these are in fact points), let $\gamma \sim \gamma'$ if $\tilde{\gamma}(1) = \tilde{\gamma}'(1)$ and $[\gamma\bar{\gamma}'] \in H$.

First we define an equivalence class as $[\gamma] \sim [\gamma']$ if and only if $[\gamma\bar{\gamma}'] \in H$. We omit the proof of reflexivity, transitivity and symmetry as it is easy to see.

We then take our universal cover \tilde{X} and identify points that belong in this equivalence class. Observe that this

For $[\gamma] = [c \cdot \gamma] \in H$ (where $[c]$ is the equivalence class of the identity), then $[c] \sim [\gamma]$. □

Another way to classify **disconnected** covering spaces of a space X is by permutations of the fiber $p^{-1}(x_0)$ for a suitable basepoint. We motivate this with a few specific examples before deriving a general construction similar to the one used to build the universal cover.




Figure 8: The three different 3 component covering spaces of the circle

Note. For some space X and baspoint x_0 , consider the 3 component covering spaces of S^1 , denoted $\tilde{X}_1, \tilde{X}_2, \tilde{X}_3$, where the subscript indicates the number of components. Let $p^{-1}(x_0)$ be the fiber of p and notice it is just three points for all \tilde{X}_n . For each covering space, the lift of the generator of $\pi_1(S^1)$ induces a permutation of the fiber. There are three starting points in $p^{-1}(x_0)$ it may start and the end of the lift is another point in $p^{-1}(x_0)$, potentially the same point.

Notice the types of permutations p induces is different for each \tilde{X}_n . \tilde{X}_3 induces cyclical permutations on all three points. \tilde{X}_2 induces cyclical permutations on two points with the third fixed. \tilde{X}_1 induces only the identity permutation.

In fact, disconnected covering spaces of S^1 are efficient graphical representations of the different permutations of the fiber $p^{-1}(x)$. This is also true for $S^1 \vee S^1$ (and perhaps true for all such spaces where the fundamental group is free on some generators).

This is no longer true if the covering space must be connected. For example, with the 3 component covering space of S^1 , the generator must lift to some permutation of all basepoints otherwise the space is disconnected.

We can generalize this idea that "covering spaces are just permutations" with the following idea. Consider $\gamma \in \pi_1(X, x_0)$, and notice each *gan̄ma* (there can be many) starts at some $p^{-1}(\gamma(0))$ and this choice of starting point defines where it ends in $p^{-1}(\gamma(1))$. We can define a bijective function $L_\gamma : p^{-1}(\gamma(0)) \rightarrow p^{-1}(\gamma(1))$.

In fact, each \tilde{X} defines a unique set of L_γ that correspond to each $[\gamma] \in \pi_1(X)$. All we have done is formalized the idea that we can construct many unique covering spaces from the same space using different ways of connecting the points in the fiber.

One important modification is to make composition of L_γ respect the direction of path composition. $\gamma \cdot \eta$ goes left to right, but $L_{\gamma\eta} = L_\gamma \cdot L_\eta$ is the opposite direction. If we define L_γ as $p^{-1}\gamma(1) \rightarrow p^{-1}\gamma(0)$ instead, composition of functions respects composition of paths.

The n -sheeted covering spaces of X are then described by the homomorphisms $\pi_1(X, x_0) \rightarrow \text{Sym}(n)$ defined as $\gamma \mapsto L_\gamma$. Here $\text{Sym}(n)$ is the permutation group of n items.

For a given covering space $\tilde{X} \rightarrow X$, the set of isomorphisms, $\tilde{X} \rightarrow \tilde{X}$, actually have a name and themselves form a group $G(\tilde{X})$. We call these **deck transformations**.

Note that each deck transformation is uniquely determined by where it maps a single point, so we will often work with basepoints without losing information.

We call a covering space **normal** if for each point $x \in X$, there exists a deck transformation between each pair of lifted points $\tilde{x}, \tilde{x}' \in \tilde{X}$. We will now show a correspondence between this geometric concept and the familiar algebraic property.

Definition 1.42. Let $p : \tilde{X} \rightarrow X$ be a covering space. Let $H = p_*(\pi_1(\tilde{X}, \tilde{x}_0))$ be a subgroup of $\pi_1(X, x_0)$.

- \tilde{X} is a normal covering space iff $H \trianglelefteq \pi_1(X, x_0)$

Exercise 1.19. Let Y be a subset of \mathbb{R}^2 that consists of a portion of the function $\sin(\frac{1}{x})$, the segment $L = [-1, 1]$ in the y -axis and an arc connecting these pieces. Let $f : Y \rightarrow S^1$ be the quotient map collapsing the segment to a single point. Show that f does not lift to the covering space $\mathbb{R} \rightarrow S^1$ despite $\pi_1(Y) = 0$. (Then $f_*(\pi_1(Y)) \subseteq p_*(\pi_1(\tilde{X}))$ would be trivial and we should have our lift be Proposition 1.36). Indeed local path connectedness is a necessary property for the lifting criterion.

Proof. The problematic component is L . Assume we have a lift \tilde{f} . We first observe that $\tilde{f}(L)$ must be a single point. By definition of the quotient map f assume $f(L) = \{1\}$ ($1 = e^{0i}$ or the top point of S^1 and the choice of point does not matter) and see $p^{-1}(\{1\})$ is a disjoint set of points in \mathbb{R} . Because L is connected, $\tilde{f}(L)$ must also be a single point.

Now see that $\tilde{f}(Y/L)$ must lie in a single component of $p^{-1}f(Y/L)$ because Y/L is connected. f is surjective and $f(Y/L)$ is $S^1 - \{0\}$, so $\tilde{f} = p^{-1}f(Y/L) = (0, 2\pi)$. However, because Y is compact, $\tilde{f}(Y)$ must also be compact and $[0, 2\pi] \subseteq \tilde{f}(Y)$ (certainly $(0, 2\pi) \neq \tilde{f}(Y)$). (It is important to see that this would not be true if L was already collapsed to a point in Y . See the note below). Then $\{0, 2\pi\} \in \tilde{f}(L)$, a contradiction. \square

Note. While $[-1, 1] \cup \{(x, \frac{1}{x}) | x \in (0, 1]\}$, the topologist's sin curve, is compact, $[-1, 1] \cup \{(x, \frac{1}{x}) | x \in (0, 1]\}$, the topologist's sin curve with the leftmost edge collapsed to a point, is not compact.

We will use some results about homotopy inverses from Chapter 0.

Exercise 1.20. Show that for $f : X \rightarrow Y$ and $g, h : Y \rightarrow X$, f is a homotopy equivalence if $fg \simeq 1_Y$ and $hf \simeq 1_X$.

More generally, if fg and hf are homotopy equivalences, f is also a homotopy equivalence.

Proof. The map hfg satisfies the required homotopy equivalence properties - $f \cdot hfg \simeq 1_X$ and $hfg \cdot f \simeq 1_Y$. To see this, $f hfg \simeq f 1_X g \simeq fg \simeq 1_Y$. A symmetric argument is used for the second equivalence.


There is also a more straightforward proof for the first part. $g \simeq gfh \simeq h$ so certainly $gf \simeq 1_X$.

To see the second result, we can use just fg . $fgi \simeq 1_Y$, so gi is a candidate for the inverse. To see $gif \simeq 1_X$, observe $ifg \simeq 1_Y$ by definition, so by left composition of g , $gifg \simeq g$ so $gif \simeq 1_X$. \square

Notice that we can swap out functions in compositions and preserve equality up to homotopy.

Exercise 1.21. Let \tilde{X} and \tilde{Y} be simply-connected covering spaces of the path-connected, locally path-connected spaces X and Y . Show that if $X \simeq Y$ then $\tilde{X} \simeq \tilde{Y}$.

Proof. Define p_X and p_Y using the commutative diagram below:



First see f and g have lifts as the necessary criteria from the existence theorem are satisfied. X and Y are both path-connected and locally path-connected, and certainly $f_*(\pi_1(X)) \subseteq p_{Y*}(\pi_1(\tilde{Y}))$ as \tilde{X} is simply connected (a symmetric claim holds for Y).

Observe $gf p_X \simeq 1_X p_X \simeq p_X$. Using the existence of lifts for g and f , we can expand the maps to arrive at $p_X \tilde{g} p_Y \tilde{f} p_X$. We can then regroup our maps to arrive at the desired relation $p_X \tilde{g} p_Y \tilde{f} p_X \simeq p_X(1_{\tilde{X}})$. Indeed, $\tilde{g} p_Y \tilde{f} p_X \simeq 1_{\tilde{X}}$. Repeating this procedure for $f g p_Y \simeq p_Y$ we recover the relation $\tilde{f} p_X \tilde{g} p_Y \simeq 1_{\tilde{Y}}$. So $p_X \tilde{f} : \tilde{X} \rightarrow \tilde{Y}$ and $p_Y \tilde{g} : \tilde{Y} \rightarrow \tilde{X}$ are homotopy inverses. \square

Exercise 1.22. Find all the 2-sheeted and 3-sheeted connected covering spaces of $S^1 \vee S^1$, up to isomorphism of the covering spaces without basepoints.

Proof. For the 2-sheeted covering spaces, we will try to use the different permutations of the fiber $p^{-1}(x)$ to identify possibilities. Let $Sym(2)$ be the permutation group of order 2 that contains just the identity map and transposition. Because $\pi_1(S^1 \vee S^1) \cong a \star b$, we can construct homomorphisms into $Sym(2)$ by mapping each of the generators, a and b onto \square

2 Appendix

2.1 Geomtry

Practice with timeless objects that are the study of algebraic topology.

Note (Real projective plane). $\mathbb{R}P^2$ Awesome explanation. <https://www.youtube.com/watch?v=B6vEdfk7SWU>

Note (Genus- g surfaces). Denoted Σ_g . g tori strung together.

2.2 Point Set Topology

Definition 2.1 (Basis). A basis for a topology \mathcal{T} on X is a collection $\mathcal{B} \subseteq \mathcal{T}$ such that every open set of the topology can be represented as a union of elements in \mathcal{B} .

Equivalently, for each $U \in \mathcal{T}$ and $x \in U$, \mathcal{B} is the collection of B where $x \in B \subseteq U$.

It is straightforward to show these definitions are equivalent (check).

Bases then admit two properties:

- \mathcal{B} covers X (X is certainly an open set that can be represented as a union of basis elements)
- For each $x \in X$ and neighborhoods $B_1, B_2 \in \mathcal{B}$, there exists $B_3 \in \mathcal{B}$ where $x \in B_3 \subseteq B_1 \cap B_2$ (As finite intersections of open sets are open, we need basis elements to fit inside each intersection)

2.3 Presentations

After more exposure to presentations to describe the group structure of some of these geometric objects, it became clear that my understanding lacked rigor.

$G = \langle X \mid R \rangle$ can be considered as the largest possible group with elements that are words composed of the letters in X **but** also subject to the relations described in R . "Subject to the relations" means that we can swap subwords for equivalent subwords or cancel subwords towards reducing different words to the same underlying group element.

(For $\langle a \mid a^3 \rangle$, such reductions look like $aaaaaa = aaa = 1$, $(a^{-1}a^{-1})a = (a)a$)

I have found this hard to reason about the equivalence of words for presentations (in abstract, not for particular groups) and it turns out that this problem is very hard. Deciding that words are equivalent in G just from the generators and relations is unsolvable in general.

Note. This problem is called the **word problem for groups**. In 1911, Max Dehn proposed that this problem was an important area of study for group theory, whereas prior most mathematicians used normal forms (irreducible representations) for group computations which made the word problem less relevant.

This does not mean that there do not exist groups where equivalences between words are obvious. Consider again $\langle a \mid a^3 \rangle$, any string / word, eg. $a^4a^{-2}a^3 \dots$, can be reduced monotonically to one of a, aa, aaa .

However, while this idea is intuitive, it is not satisfying, and a more formal definition of the group describe by a presentation is desired:

$$G = \langle X \mid R \rangle = F(X) / \langle R \rangle^{F(X)}$$

Definition 2.2. The normal closure of H in G is the smallest normal subgroup containing H . It is computed by $\langle \{xHx^{-1} \mid x \in G\} \rangle$

We will prove three properties of the presentation using this formal definition [motivated by this post](#).

Proposition 2.3. G is generated by the images of X in the quotient group.

Proof. Let $N = \langle R \rangle^{F(X)}$. Clearly the generating set $\{xN \mid x \in X\} \subseteq F(X)/N$. Then the generated set $\langle \{xN \mid x \in X\} \rangle \subseteq F(X)/N$ as any word $xNyN \dots zN$ can be reduced to $(xy \dots z)N$ and this word is in $F(X)/N$.

To see the reverse, consider any $gN \in F(X)/N$ and observe g is some word of elements of X . Expand g to the equivalent representation in letters of X , $(xy \dots z)N$ and see this element lies in the desired generated set.

We finish this proof by constructing an isomorphism by mapping each coset of $\langle \{xN \mid x \in X\} \rangle$ to the coset in that shares its representative $\subseteq F(X)/N$. \square

Proposition 2.4. G is generated by the images of X in the quotient group.

Note (Free Groups). See the [myasnikov.1.free.groups.pdf](#) for a category theory flavored exposition on free groups, their universal property and role in presentations.

Note (Universal Property of Free Groups). Let S be the generating set of a group $F(S)$. For any group G and associated map $f : S \rightarrow G$ (note f is not a homomorphism and is just mapping symbols into a group), f extends to a unique (homomorphism) $f' : F(S) \rightarrow G$ that commutes in the following diagram.

$$\begin{array}{ccc} S & \xrightarrow{f} & G \\ i \downarrow & \nearrow f' & \\ F(S) & & \end{array}$$

We proceed with a brief proof and then an example.

If we consider $F(S)$ as reduced words in letters of S , let $f' = f(s_1) \dots f(s_n)$ and claim this is our extended homomorphism.

Uniqueness can be seen because f' must agree with f for each $s \in S$.

Lets construct two isomomorphisms between presentations and groups to understand what a quotient of this enormous free group and almost just as enormous normal subgroup actually mean in the context of groups we already understand.

- $\mathbb{Z} \times \mathbb{Z} \cong \langle a, b \mid [a, b] \rangle$
- $\mathfrak{S}_3 \cong \langle a, b \mid a^2 = b^3 = aba^{-1}b^2 \rangle$

Proposition 2.5. $D_3 \cong \langle x, y \mid x^3 = y^2 = xyx^{-2}y \rangle$

Proof. Lets begin by invoking the universal property:

$$\begin{array}{ccc} \{x, y\} & \xrightarrow{i} & F(x, y) \\ & \searrow f & \downarrow \phi \\ & & D_3 \end{array}$$

Where $F(x, y)$ is our free group on two generator, i is the inclusion map, f is a map from our generators to letters in the dihedral group. The diagram commutes ($f = \phi i$).

By definition, our presentation is $F(x, y) / \langle\langle x^2, y^3, xyx^{-2}y \rangle\rangle$ (where the $\langle\langle - \rangle\rangle$ denotes the normal closure). I claim that our universal θ gives us the desired isomorphism when restricted to the quotient group.

Let us expand our commutative diagram by factoring ϕ through the quotient group:

$$\begin{array}{ccc} \{x, y\} & \xrightarrow{i} & F(x, y) \\ & \searrow f & \downarrow \pi \\ & & D_3/N \\ & & \downarrow \phi' \\ & & D_3 \end{array}$$

Where $\pi = w \rightarrow w \langle\langle x^2, y^3, xyx^{-2}y \rangle\rangle$ projects words onto their cosets and $\phi' = wN \rightarrow \phi(w)$. Notice ϕ' is well defined as the image of every element of the normal closure is the identity (" ϕ respects the relations").

Then ϕ' is surjective as ϕ is surjective (every element of D_3 is word of letters in the image of f). To see ϕ' is injective, consider distinct cosets $aN \neq bN$. Then $\phi'(aN) = \phi'(a)\phi'(N) = \phi'(a) = \phi(a)$. Similarly, $\phi'(bN) = \phi(b)$. $\phi(a) \neq \phi(b)$

□

Note. Tietze transforms define operations on presentations that do not change the generated group. Adding and removing both generators and relations are allowed if the generator or relation can be derived from other information in the presentation. $\langle x, y \mid x^2 = y = 1 \rangle$

2.4 Derived Subgroups

Commutators and derived subgroups appear frequently to describe the structure of fundamental groups of cell complexes. We will review some definitions and proofs from elementary algebra.

Definition 2.6. The **commutator** of two elements $a, b \in G$ is $a^{-1}b^{-1}ab$ and is denoted $[a, b]$.

Definition 2.7. a^x is the conjugate of a by x ($xa x^{-1}$).

Clearly the commutator is the identity element if a and b commute in G (hence its name). We derive some pedagogical identities that will be useful.

Definition 2.8. $[a, b]^x = [a^x, b^x]$

Proof. $xa^{-1}b^{-1}abx^{-1} = xa^{-1}x^{-1}xb^{-1}x^{-1}xax^{-1}xbx^{-1} = [a^x, b^x]$ □

Note. The set of commutators need not be closed over the group operation.

Consider the free group on four generators. Then $[a, b][c, d] = a^{-1}b^{-1}abc^{-1}d^{-1}cd$. This is not a commutator in this group (there are no two elements x, y such that $[x, y] = [a, b][c, d]$).

Even though products of commutators are not themselves commutators in general, the subgroup *generated* by all commutators in a group is certainly closed over the group operation. This leads naturally to the derived subgroup.

Definition 2.9. The group generated by all of the commutators in G is the **derived subgroup** of G , denoted $[G, G]$ or G' .

Theorem 2.10. $[G, G] \trianglelefteq G$

Proof. $([a, b] \cdots [e, f])^x = ([a^x, b^x] \cdots [e^x, f^x])$ □

Theorem 2.11. *If the quotient group G/N is abelian, then $[G, G] \subseteq N$. In other words, the derived subgroup is the smallest subgroup that abelianizes G .*

Proof. Let N be some normal subgroup of G , then N abelianizes G iff for each $x, y \in G$, $xyN = yxN$. Then $N = y^{-1}x^{-1}yxN$ (the group operation is well-defined as N is normal) so $y^{-1}x^{-1}yx \in N$.

Certainly the set of commutators must be included in N . So any minimal N is the minimal group generated by these commutators. But this is exactly $[G, G]$. □

Definition 2.12. $G/[G, G]$ is the abelianization of G and sometimes denoted G^{ab} .

2.5 Continuity

I found myself confused about the equivalence between the $\delta\epsilon$ definition of continuity from calculus and the topological definition using open sets. The direction of the implications is what is not obvious.

Recall the conceptual meat of continuity is that small perturbations in the range can always be .

More specifically the preimage of some small interval in the range always exists in the domain if we

Definition 2.13. A function is continuous at x if for all ϵ there exists an δ where

$$|x - x_0| < \delta \implies |f(x) - f(x_0)| < \epsilon$$

The structure of this implication is crucial. We can construct arbitrarily small intervals of the range and always recover some δ about x_0 that entirely maps within this interval. Simple piecewise discontinuous functions from analysis illustrate how clearly discontinuous maps (they have big holes) map onto this delta epsilon definition. We will walk through one for the sake of pedagogy.

Example 2.14. Consider

$$\begin{cases} x^2 - 3 & x \leq 0 \\ \sin x & x > 0 \end{cases}$$

Consider continuity at $x = 0$. For $\epsilon = 2$, there exists no value of δ where $|x - 0| \implies |f(x) - f(0)| < 2$.

Continuity is defined differently in topology:

Definition 2.15. Let $f : X \rightarrow Y$ be a continuous map. Then for each open $U \in Y$, $f^{-1}(U)$ is open in X .

2.6 Complex Numbers

$z = a + bi$ can be represented as

$$e^{i\theta} = \sin \theta + i \cos \theta$$